

Ring-necked Pheasant Status Report, 2017

Agriculture and Grassland Wildlife Program Note 17-3



Ring-necked Pheasant in Illinois

Pheasant hunting in Illinois was very popular in the 1960's and 70's when 250,000 hunters harvested over a million birds per year several times. Due primarily to changes in land use and farming practices, this game bird that had been such an abundant by-product of the agricultural landscape began a dramatic decline in the mid 1970's (Figure 1). By the year 2000, pheasant hunters had declined to 59,000 harvesting only 158,000 birds. The small, diverse farms of the 1950's had abundant small grains (wheat, oats etc.), livestock, and hay fields with lots of fencerows and hedgerows separating small fields. Today, two crops (corn and soybeans) have replaced most of the small grains, pastures, and hayfields so common in the 1960's and 70's. The remaining pastures and waterways are mostly planted to cool-season grasses like brome and fescue that get too thick to provide quality habitat for pheasants or quail. Hedgerows and fencerows separated small fields and provided cover for game are sparse or have been 'cleaned up' with herbicide.

Field size and the size of equipment have also increased dramatically in the last 50 years. Larger and more efficient equipment decrease the amount of time it takes to plant and harvest crops. Most fields are disked soon after harvest, covering most crop stubble early in the fall. There is very little cover left for wildlife over the winter and crops get planted quickly in the spring. During the summer, 'recreational' or aesthetic mowing of roadsides and non-crop areas often occurs during prime nesting season (April – August) for the ring-necked pheasant and other grassland birds. The changes to the landscape and the loss of habitat have occurred slowly, over several decades, but the additive effects on pheasant and quail populations are dramatic.




Figure 1 – Ring-necked Pheasant Harvest and Pheasant Hunters 1955 – 2016.

2016-17 Hunter Harvest

Harvest estimates in this report are based on results of the 2016-17 Illinois Hunter Harvest Survey prepared annually by the Illinois Natural History Survey. During the 2016-17 pheasant season, an estimated 10,204 hunters (26% decrease from last year) harvested 14,831 wild pheasant (40% decrease). Even though the number of hunters decreased, the number of days each hunter spent in the field also decreased from 2015.






Figure 2 – Pheasant Harvest and Hunters 2011 - 2016

Pheasant Population Index

There are established routes across the state and biologists run the routes two times annually during the breeding season. The first run is completed during the peak breeding window for ring-necked pheasant (May 10 – June 10). Each route has 20 stops for 3 minutes each to record individuals heard or seen. The number of routes surveyed annually depends on staff availability, but as many of the surveys are completed as possible.

During 2017, observers recorded an average of 0.26 pheasants per stop on the 71 survey routes (30% decrease). Ring-necked pheasants were recorded at 14% of the stops (percent occurrence) on the routes (33% decrease). The number of pheasants counted and the number of stops where pheasants were seen or heard were lower than the numbers from the 2016 surveys.



Figures 3 and 4 – Birds per Stop and Percent Occurrence for Pheasant on Upland Routes Weather 2016-17

The winter of 2016-17 was relatively mild and winter mortality due to weather was likely low for pheasant. Spring brought above-average rainfall and may have been problematic for nesting hens and broods in May and early June. From late June through most of August, weather conditions improved across much of the state (with parts of northern Illinois being the exception due to record flooding). Overall, 2017 weather conditions were a little wet for pheasant during their peak nesting season, but good for hens that attempted a second nest.

Despite some heavy rains early, a recent pheasant study conducted in east-central Illinois highlights the importance of having high-quality habitat to encourage recruitment into the pheasant population. Despite some inclement weather, nesting effort, nest success and brood survival on some of the state-owned Pheasant Habitat Areas and other, high-quality private grasslands has been good in years with similar weather. The research on these sites found the areas hen pheasants seek for nesting and raising their broods are often the same areas used to take cover during heavy snow events. The presence of thick-stemmed and hardy forbs keeps grasses upright during the winter when winds, rains and snow can often lay solid stands of native grasses flat.

2017-18 Season Outlook

The key to a successful 2017-18 pheasant season is finding high-quality habitat. Focus on areas with relatively new CRP plantings, or recently managed grasslands, with hay and/or small grains nearby. Within these areas, look for abundant forbs (broad-leafed, flowering plants) mixed in with scattered clumps of grasses. Hunters may find fewer roosters across Illinois this fall based on lower survey numbers this summer.

Management Tips

- An important management tip to benefit pheasant (as well as many other grassland birds) is to break up mowing times to leave some quality habitat throughout the year. Do not mow more than 1/3 or 1/4 of any field, waterway or fallow area at any one time. This strategy is also important to provide nectar sources for pollinators like monarch butterflies, as well as habitat for grassland birds and other wildlife from early spring to late fall.
- Avoid mowing during peak nesting season, from April 15 – August 1. Mowing during the nesting season can destroy nests and often kills the hen as well.
- It is important to ensure there will always be quality nesting, brood-rearing and escape cover in your grassland. This means only managing a portion (1/3 or 1/4) of your habitat at any one time and planning ahead for which types of cover you are managing for on each part of your grassland.
- Fall mowing or burning in September or early October can clear away thatch and stimulate vigorous regrowth of cool-season grasses that invade stands of native grasses. Once this new growth reaches 6” – 12” in the fall, read and follow label directions for a foliar application of herbicide like glyphosate to set back areas of tall fescue, brome, bluegrass or reed-canary grass. A second herbicide application may be needed the following spring/summer before native grasses begin growing.
- Early fall is also a good time to spray and/or disc small blocks in existing grasslands that have become too thick. These blocks should provide good brood habitat the following summer.

Thank you for your interest in the ring-necked pheasant. Have a safe and enjoyable hunting season in 2017!

Season Dates

November 4, 2017 – January 8, 2018 (North)
November 4, 2017 – January 15, 2018 (South)

Hours

Sunrise to Sunset

Bag Limits

2 per day with 6 in possession after the 3rd day

Upland Game, Furbearers & Turkey Zones

Upland Game:
Cock Pheasant, Quail, Hungarian Partridge, Rabbit: Divided by U.S. Route 36 from the Indiana state line to Springfield, Illinois Route 29 from Springfield to Pekin, and Illinois Route 9 from Pekin to Dallas City, then due west to the Mississippi River.

Furbearers:
hunting (Raccoon, Opossum) and trapping: Divided by U.S. Route 36.

Turkey:
North Zone includes that portion of Illinois north of Crawford, Jasper, Effingham, Fayette, Bond and Madison counties. South Zone includes the remainder of the state.

Legend:
- - - - - Upland Game
———— Furbearers
· · · · · Turkey



photo by Adele Hodde

*Prepared by Stan McTaggart 10/17
Illinois Department of Natural Resources
Division of Wildlife Resources*

Equal opportunity to participate in programs of the Illinois Department of Natural Resources (IDNR) and those funded by the U.S. Fish and Wildlife Service and other agencies is available to all individuals regardless of race, sex, national origin, disability, age, religion or other non-merit factors. If you believe you have been discriminated against, contact the funding source's civil rights office and/or the Equal Employment Opportunity Officer, IDNR, One Natural Resources Way, Springfield, IL 62702-1271; 217/785-0067; TTY 217/782-9175.